



Generalization of matching extensions in graphs (III)

Bing Bai^a, Hongliang Lu^b, Qinglin Yu^{c,d,*}

^a Center for Combinatorics, LPMC, Nankai University, Tianjin, PR China

^b Department of Mathematics, Xian Jiaotong University, Xian, PR China

^c Department of Mathematics and Statistics, Thompson Rivers University, Kamloops, BC, Canada

^d School of Mathematics, Shandong University, Jinan, Shandong, PR China

ARTICLE INFO

Article history:

Received 24 May 2010

Received in revised form 22 December 2010

Accepted 29 December 2010

Available online 22 January 2011

Keywords:

(n, k, d) -graphs

k -extendable graphs

n -factor-critical graphs

ABSTRACT

Proposing them as a general framework, Liu and Yu (2001) [6] introduced (n, k, d) -graphs to unify the concepts of deficiency of matchings, n -factor-criticality and k -extendability. Let G be a graph and let n, k and d be non-negative integers such that $n + 2k + d + 2 \leq |V(G)|$ and $|V(G)| - n - d$ is even. If on deleting any n vertices from G the remaining subgraph H of G contains a k -matching and each k -matching can be extended to a defect- d matching in H , then G is called an (n, k, d) -graph. In this paper, we obtain more properties of (n, k, d) -graphs, in particular the recursive relations of (n, k, d) -graphs for distinct parameters n, k and d . Moreover, we provide a characterization for maximal non- (n, k, d) -graphs.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

All graphs considered in this paper are finite, connected, loopless and have no multiple edges. For the most part our notation and terminology follow those of Bondy and Murty [3].

Let G be a graph with vertex set $V(G)$, edge set $E(G)$ and minimum degree $\delta(G)$. A *matching* M of G is a subset of $E(G)$ such that any two edges of M have no vertices in common. A matching of k edges is called a k -*matching*. For a matching M , we use $V(M)$ to denote the vertices incident to the edges of M . Let d be a non-negative integer. A matching is called a *defect- d matching* if it covers exactly $|V(G)| - d$ vertices of G . Clearly, a defect-0 matching is a perfect matching. For a subset S of $V(G)$, we denote by $G[S]$ the subgraph of G induced by S and we write $G - S$ for $G[V(G) \setminus S]$. The number of odd components of G is denoted by $c_0(G)$. The *join* $G \vee H$ of two graphs G and H is a graph with vertex set $V(G) \cup V(H)$ and edge set $E(G) \cup E(H) \cup \{xy : x \in V(G), y \in V(H)\}$. We denote the complement of G by \bar{G} . A set T is called an n -*set* if $|T| = n$. For two disjoint sets A and B of $V(G)$, we define $E(A, B) = \{xy : x \in A \text{ and } y \in B\} \cap E(G)$.

Let M be a matching of G . If there is a matching M' of G such that $M \subseteq M'$, we say that M can be extended to M' or M' is an *extension* of M . Suppose that G is a connected graph with perfect matchings. If each k -matching can be extended to a perfect matching in G , then G is called *k-extendable*. To avoid triviality, we require that $|V(G)| \geq 2k + 2$ for k -extendable graphs. This family of graphs was introduced by Plummer [9]. A graph G is called *n-factor-critical* if after deleting any n vertices the remaining subgraph of G has a perfect matching. This concept is introduced by Favaron [4] and Yu [10], independently, and is a generalization of the notions of the well-known factor-critical graphs and bicritical graphs, the cases of $n = 1$ and 2, respectively. In [8], Lou investigated the relationship between $2k$ -factor-criticality and k -extendability.

Let G be a graph and let n, k and d be non-negative integers such that $|V(G)| \geq n + 2k + d + 2$ and $|V(G)| - n - d$ is even. If on deleting any n vertices from G the remaining subgraph of G contains a k -matching and each k -matching in the subgraph can be extended to a defect- d matching, then G is called an (n, k, d) -*graph*. This term was introduced by Liu and Yu [6] as

* Corresponding author at: Department of Mathematics and Statistics, Thompson Rivers University, Kamloops, BC, Canada. Tel.: +1 250 371 5552.
E-mail address: yu@tru.ca (Q. Yu).

a general framework for unifying the concepts of defect- d matchings, n -factor-criticality and k -extendability. In particular, $(n, 0, 0)$ -graphs are exactly n -factor-critical graphs and $(0, k, 0)$ -graphs are just the same as k -extendable graphs. In [5,6], the recursive relations were shown for distinct parameters n , k and d and the impact of adding or deleting an edge for $d \geq 0$ was discussed. In this paper, we continue the investigation of (n, k, d) -graphs and obtain more recursive relations.

A graph G is called a *maximal non- (n, k, d) -graph* if G is not an (n, k, d) -graph but $G \cup e$ is an (n, k, d) -graph for every edge $e \in E(G)$. In [1], Ananchuen et al. studied maximal non- k -factor-critical graphs and maximal non- k -extendable graphs; they also provided a characterization of these graphs. In the current paper, we generalize their criteria to obtain a characterization of maximal non- (n, k, d) -graphs.

2. Known results

A necessary and sufficient condition for a graph to have a defect- d matching was given by Berge [2].

Lemma 2.1 (Berge [2]). *Let G be a graph and d an integer such that $0 \leq d \leq |V(G)|$ and $|V(G)| \equiv d \pmod{2}$. Then G has a defect- d matching if and only if for any $S \subseteq V(G)$*

$$c_0(G - S) \leq |S| + d.$$

In [6], Liu and Yu showed the following sufficient and necessary conditions for (n, k, d) -graphs.

Lemma 2.2 (Liu and Yu [6]). *A graph G is an (n, k, d) -graph if and only if the following conditions hold:*

(a) *for any $S \subseteq V(G)$ such that $|S| \geq n$, then*

$$c_0(G - S) \leq |S| - n + d,$$

(b) *for any $S \subseteq V(G)$ such that $|S| \geq n + 2k$ and $G[S]$ contains a k -matching, then*

$$c_0(G - S) \leq |S| - n - 2k + d.$$

It is a natural to try to find recursive relations among the graphs with different parameters n , k and d . Below is one result of such an endeavor.

Lemma 2.3 (Liu and Yu [6]). *Every (n, k, d) -graph is also an (n', k', d) -graph, where $0 \leq n' \leq n$, $0 \leq k' \leq k$ and $n' \equiv n \pmod{2}$.*

3. Main results

Following the study of recursive relations of the previous work, we continue to investigate the effect of various graphic operations on (n, k, d) -graphs and recursive relations. We start with the following lemma.

Lemma 3.1. *If G is an (n, k, d) -graph, then it is also an $(n - 2, k + 1, d)$ -graph.*

Proof. First, note that G is an $(n - 2, 0, d)$ -graph by Lemma 2.3. Since $|V(G)| \geq n + 2k + d + 2$, for any $(n - 2)$ -set $S \subseteq V(G)$ there exist $(k + 1)$ -matchings in subgraph $G - S$.

Suppose, to the contrary, that G is not an $(n - 2, k + 1, d)$ -graph. Then, by the definition, there exist an $(n - 2)$ -set $R \subseteq V(G)$ and a $(k + 1)$ -matching M which cannot be extended to a defect- d matching of $G - R$. By Lemma 2.1 and parity, there exists a subset S_0 in $G - R - V(M)$ such that

$$c_0(G - R - V(M) - S_0) \geq |S_0| + d + 2.$$

Let $S = S_0 \cup R \cup V(M)$. Then $|S| = |S_0| + |R| + 2(k + 1) \geq n + 2k$ and $G[S]$ contains k -matchings, and

$$c_0(G - S) = c_0(G - S_0 - R - V(M)) \geq |S_0| + d + 2 = |S| - n - 2k + d + 2,$$

a contradiction to Lemma 2.2(b). \square

Theorem 3.2. *A graph G is an $(n + 2, k - 1, d)$ -graph if and only if G is an (n, k, d) -graph and $G \cup e$ is an (n, k, d) -graph, for any $e \in E(\bar{G})$.*

Proof. If G is an $(n + 2, k - 1, d)$ -graph, by Lemma 3.1, then G is an (n, k, d) -graph.

We show that $G \cup e$ is an (n, k, d) -graph for any $e \in E(\bar{G})$. Otherwise, there exists an edge $e_1 \in E(\bar{G})$ such that $G' = G \cup \{e_1\}$ is not an (n, k, d) -graph. By Lemma 2.2, we consider two cases:

Case 1. There exists a subset $S_1 \subseteq V(G') = V(G)$ such that $|S_1| \geq n$ and $c_0(G' - S_1) \geq |S_1| - n + d + 2$. However,

$$c_0(G - S_1) \geq c_0(G' - S_1) \geq |S_1| - n + d + 2,$$

a contradiction to G being an (n, k, d) -graph and Lemma 2.2(a).

Case 2. There exists a subset $S_2 \subseteq V(G') = V(G)$, where $|S_2| \geq n + 2k$ and $G'[S_2]$ contains a k -matching M_2 such that

$$c_0(G' - S_2) \geq |S_2| - n - 2k + d + 2.$$

If $e_1 \notin M_2$, then $|S_2| \geq n + 2k$ and $G[S_2]$ contains the k -matching M_2 , and $c_0(G - S_2) \geq c_0(G' - S_2) \geq |S_2| - n - 2k + d + 2$, a contradiction to G being an (n, k, d) -graph and Lemma 2.2(b). So $e_1 \in M_2$. Let $M'_2 = M_2 - \{e_1\}$. Then $|S_2| \geq n + 2k = (n + 2) + 2(k - 1)$ and $G[S_2]$ contains the $(k - 1)$ -matching M'_2 . Moreover,

$$c_0(G - S_2) \geq c_0(G' - S_2) \geq |S_2| - n - 2k + d + 2 = |S_2| - (n + 2) - 2(k - 1) + d + 2,$$

a contradiction to G being an $(n + 2, k - 1, d)$ -graph.

Next we prove the sufficiency. Suppose that G is not an $(n + 2, k - 1, d)$ -graph. Then there exist an $(n + 2)$ -set $S_3 \subseteq V(G)$ and a $(k - 1)$ -matching M_3 which cannot be extended to a defect- d matching of $G - S_3 - V(M_3)$. By Lemma 2.1, there exists a vertex set $R \subseteq V(G - S_3 - V(M_3))$ such that

$$c_0(G - S_3 - V(M_3) - R) \geq |R| + d + 2.$$

For any two vertices u, v of S_3 , if $uv \in E(\bar{G})$, let $e_2 = uv$, $M'_3 = M_3 \cup \{e_2\}$, and $S'_3 = S_3 \setminus \{u, v\}$; then we have

$$c_0((G \cup e_2) - S'_3 - V(M'_3) - R) = c_0(G - S_3 - V(M_3) - R) \geq |R| + d + 2,$$

a contradiction to the fact that $G \cup e$ is an (n, k, d) -graph, for any $e \in E(\bar{G})$; if $uv \in E(G)$, then $|S'_3| = n$ and M'_3 is a k -matching of G , and

$$c_0(G - S'_3 - V(M'_3) - R) = c_0(G - S_3 - V(M_3) - R) \geq |R| + d + 2,$$

a contradiction to G being an (n, k, d) -graph. \square

Applying Lemma 3.1, we have sufficient and necessary conditions for $(n + 2k, 0, d)$ -graphs.

Theorem 3.3. A graph G is an $(n + 2k, 0, d)$ -graph if and only if G is an (n, k, d) -graph and for any edge set $D \subseteq E(\bar{G})$, $G \cup D$ is an (n, k, d) -graph.

Proof. If G is an $(n + 2k, 0, d)$ -graph, clearly $G \cup D$ is also an $(n + 2k, 0, d)$ -graph. Applying Lemma 3.1 repeatedly, we see that $G \cup D$ is an (n, k, d) -graph.

On the other hand, suppose that G is not an $(n + 2k, 0, d)$ -graph; by Lemma 2.2, there exists a subset S with $|S| \geq n + 2k$ such that

$$c_0(G - S) \geq |S| - (n + 2k) + d + 2.$$

Let $S = \{u_1, \dots, u_h\}$, where $h \geq n + 2k$ and $G' = G \cup \{u_{2i-1}u_{2i} \mid i = 1, \dots, k\}$. Then $G'[S]$ contains a k -matching and we have

$$c_0(G' - S) = c_0(G - S) \geq |S| - (n + 2k) + d + 2.$$

By Lemma 2.2(b), G' is not an (n, k, d) -graph, a contradiction. \square

Let $n = 0$ and $d = 0$; we have the next corollary.

Corollary 3.4 (Lou [8]). A graph G of even order is $2k$ -factor-critical if and only if:

- (a) G is k -extendable; and
- (b) for any edge set $D \subseteq E(\bar{G})$, $G \cup D$ is k -extendable.

In [7], Liu and Yu present several results concerning $(n, k, 0)$ -graphs and its subgraphs. In particular, they proved that if $G - V(e)$ is an $(n, k, 0)$ -graph for each $e \in F$ (where F is a fixed 1-factor in G), then G is an $(n, k, 0)$ -graph. We generalize this result for any $d \geq 0$ and $n \geq d + 2$.

Theorem 3.5. Let F be a perfect matching of a connected graph G , where $|V(G)| \geq n + 2k + d + 4$ and $n \geq d + 2$. If subgraph $G - V(e)$ is an (n, k, d) -graph for each $e \in F$, then G is also an (n, k, d) -graph.

Proof. Assume that F is a perfect matching of G such that $G - V(e)$ is an (n, k, d) -graph for each $e \in F$. To see the existence of k -matchings in the subgraphs, we show a claim.

Claim 1. For any n -set $T \subseteq V(G)$, $G - T$ contains k -matchings.

If $F \cap E(G - T) = \emptyset$, then there exists an edge $e = ab \in F$ such that $a \in T$ and $b \in V(G - T)$. Let $T' = T \setminus \{a\} \cup \{c\}$, where $c \in V(G) - T - \{b\}$. Then $|T'| = n$ and $F \cap E(G - T') = \{e\}$. By the assumption of the theorem, $G - V(e)$ is an (n, k, d) -graph. Hence, $G - V(e) - T'$ has a defect- d matching M_1 . Since $|V(G)| \geq n + 2k + d + 4$, M_1 contains at least $k + 1$ edges. Therefore, $G - T$ contains k -matchings.

If $F \cap E(G - T) \neq \emptyset$, let $e = ab \in F \cap E(G - T)$; then $G - V(e)$ is an (n, k, d) -graph. So $G - V(e) - T$ contains k -matchings and thus $G - T$ contains k -matchings.

Suppose that G is not an (n, k, d) -graph; by the definition and Claim 1, there exists a vertex set R of order n in G and a k -matching M of $G - R$ such that $G - R - V(M)$ has no defect- d matchings. Let $G' = G - R - V(M)$; by Lemma 2.1 and parity, there exists a subset S in G' such that

$$c_0(G' - S) = c_0(G - R - V(M) - S) \geq |S| + d + 2. \quad (1)$$

Claim 2. $F \cap E(G[R \cup S]) = F \cap M = F \cap E(V(M), R \cup S) = F \cap E(C_i) = F \cap E(S, V(C_i)) = \emptyset$ for all C_i , where C_i is an odd component of $G' - S$.

If there exists an edge $e \in (F \cap E(R)) \cup (F \cap E(S))$, say $e \in F \cap E(R)$, then we have

$$c_0(G - V(e) - (R \setminus V(e)) - V(M) - S) = c_0(G' - S) \geq |S| + d + 2.$$

So $G - V(e)$ is not an $(n - 2, k, d)$ -graph, a contradiction to $G - V(e)$ being an (n, k, d) -graph and Lemma 2.3.

If there exists an edge $e \in F \cap E(R, S)$, where $e = ab$, $a \in S$, $b \in R$, let $c \in C_i$, $R' = R \setminus \{b\} \cup \{c\}$, and $S' = S \setminus \{a\}$. Then we have

$$c_0(G - V(e) - R' - V(M) - S') \geq c_0(G' - S) - 1 \geq |S'| + d + 2.$$

Thus $G - V(e)$ is not an (n, k, d) -graph, a contradiction.

If there exists an edge $e \in F \cap M$, then we have

$$c_0(G - V(e) - R - V(M \setminus \{e\}) - S) = c_0(G' - S) \geq |S| + d + 2.$$

Thus $G - V(e)$ is not an $(n, k - 1, d)$ -graph, a contradiction.

Suppose that $e \in F \cap E(V(M), R)$. Let $e = uv$ and $ua \in M$, where $u \in V(M)$ and $v \in R$. Let $R_1 = (R \setminus \{v\}) \cup \{a\}$ and $M'' = M \setminus \{ua\}$. Then

$$c_0(G - V(e) - R_1 - V(M'') - S) \geq |S| + d + 2.$$

Thus $G - V(e)$ is not an $(n, k - 1, d)$ -graph, a contradiction.

Using the similar arguments, we may show that $e \notin E(S) \cup E(V(M), S) \cup (\cup_i E(C_i)) \cup E(S, V(C_i))$ for any $e \in F$.

Claim 3. $G' - S$ has no even components.

Otherwise, let D be an even component of $G' - S$ and $e = ab \in F$, $a \in V(D)$. If $b \in R$, choose a vertex $c \in V(D) \setminus \{a\}$, let $R_2 = R \setminus \{b\} \cup \{c\}$; then

$$c_0(G - V(e) - R_2 - V(M) - S) \geq c_0(G' - S) \geq |S| + d + 2.$$

Thus $G - V(e)$ is not an (n, k, d) -graph, a contradiction. For $b \in S$, we arrive at a contradiction with a similar argument. So we may assume that $b \in V(M)$. Let $bc \in M$. Set $S_1 = S \cup \{c\}$. Note that $G'[D \setminus \{a\}]$ contains at least one odd component. So we have

$$c_0(G - V(e) - R - V(M \setminus \{bc\}) - S_1) \geq |S_1| + d + 2.$$

Hence $G - V(e)$ is not an $(n, k - 1, d)$ -graph, a contradiction.

Finally, if e is in the component D , then

$$c_0(G - V(e) - R - V(M) - S) \geq c_0(G' - S) \geq |S| + d + 2.$$

Thus $G - V(e)$ is not an (n, k, d) -graph, a contradiction again.

For any vertex $x \in S$, by Claim 2 x cannot be matched in perfect matching F to any other vertex in S or any vertex in $R \cup V(M)$ or any vertex in an odd component, so we conclude that $S = \emptyset$.

Claim 4. $c_0(G' - S) = c_0(G') = d + 2$.

By (1), we need only show that $c_0(G') \leq d + 2$. Otherwise, suppose $c_0(G') \geq d + 3$. If there exists an edge $e = ab \in F \cap E(R, C_i)$, where $a \in C_i$ and $b \in R$, we choose a vertex x from another odd component C_j and let $R_1 = R \setminus \{b\} \cup \{x\}$; then

$$c_0(G - V(e) - R_1 - V(M)) \geq c_0(G') - 2 \geq d + 1.$$

Thus $G - V(e)$ is not an (n, k, d) -graph, a contradiction. Next, we assume that all vertices in $\cup_i C_i$ are matched to $V(M)$. Consider the alternating path $P = c_i x_1 y_1 \dots x_m y_m c_j$ of $F \cup M$ starting at C_i and ending at C_j . Let $e = c_i x_1 \in F$ and $M' = M \triangle (P \setminus \{e\})$. Then

$$c_0(G - V(e) - R - V(M')) \geq c_0(G') - 2 \geq d + 1,$$

a contradiction.

Now we proceed to the proof of the theorem.

Since $|V(G')| \geq d + 4$ and $c_0(G') = d + 2$, there exists one odd component of order at least 3. Moreover, as $n \geq d + 2$, $c_0(G') = d + 2$ and $F \cap (E(R, V(M)) \cup E(R)) = \emptyset$, there must exist an edge $e = ab \in F$ from R to an odd component C_i with $|C_i| \geq 3$, where $a \in C_i$ and $b \in R$. Since $|C_i| \geq 3$, choose a vertex $x \in C_i \setminus \{a\}$. Let $R_2 = R \setminus \{b\} \cup \{x\}$. Then

$$c_0(G - V(e) - R_2 - V(M)) \geq c_0(G') = d + 2,$$

a contradiction.

We complete the proof. \square

In [5], Jin et al. proved the recursive relation for adding a vertex.

Theorem 3.6 (Jin et al. [5]). *Let G be an (n, k, d) -graph with $k > 0$ and $n > d$. Then $G \vee x$ is an $(n + 1, k - 1, d)$ -graph for any vertex $x \notin V(G)$.*

Here we present an example to show that the condition $n > d$ is necessary.

For $k > 0$ and $n \leq d$, let $d = n + r$ for some $r \geq 0$. We consider a bipartite graph $H = K_{m, m+r}$, where $m \geq n + k$. Then H is an $(n, k, n + r)$ -graph, but $H \vee x$ is not an $(n + 1, k - 1, n + r)$ -graph.

4. Maximal non- (n, k, d) -graphs

In this section, we provide a characterization of maximal non- (n, k, d) -graphs, which is a generalization of the characterization of maximal non- k -factor-critical graphs in [1].

Theorem 4.1. *Let G be a connected graph of order p and n, k, d be positive integers with $p + n + d \equiv 0 \pmod{2}$. Then G is a maximal non- (n, k, d) -graph if and only if*

$$G \cong K_{n+2k+s} \vee \left(\bigcup_{i=1}^{s+d+2} K_{2t_i+1} \right),$$

where s and t_i are non-negative integers with $\sum_{i=1}^{s+d+2} t_i = \frac{p-n-2k-d}{2} - s - 1$.

Proof. Let $H = K_{n+2k+s}$ and $G_i = K_{2t_i+1}$ for $1 \leq i \leq s + d + 2$. Suppose that the theorem does not hold. That is, there exists an edge $e \in E(\bar{G})$ such that $G' = G \cup e$ is not an (n, k, d) -graph. Then e is an edge connecting G_i and G_j for some i and j .

By Lemma 2.2 and the parity argument, then either:

- (a) there exists a subset S' in G' with $|S'| \geq n$ and $c_0(G' - S') \geq |S'| - n + d + 2$; or
- (b) there exists a subset S' in G' such that $|S'| \geq n + 2k$ and S' contains a k -matching satisfying $c_0(G' - S') \geq |S'| - n - 2k + d + 2$.

Clearly, $V(H) \subseteq S'$ and so S' contains a k -matching. Thus we need only consider (b). Hence we have $c_0(G' - S') \geq |S'| - n - 2k + d + 2 \geq |V(H)| - n - 2k + d + 2 \geq d + s + 2$. If $c_0(G' - S') = d + s + 2$, then $|S'| = n + 2k + s$ and so $S' = V(H)$. Therefore we have $c_0(G' - S') = d + s$, a contradiction. Hence we have $|S'| > n + 2k + s$ and then $c_0(G' - S') > d + s + 2$. But $G' - S'$ contains at most $s + d + 2$ odd components, a contradiction.

Now we prove the necessity. Since G is a maximal non- (n, k, d) -graph, for any n -subset R of $V(G)$ there exists a k -matching M in $G - R$. Let $G' = G - R - V(M)$. By Lemma 2.1 and parity, there exists a set S' in G' such that

$$c_0(G' - S') \geq |S'| + d + 2.$$

Let C_1, C_2, \dots, C_r be odd components in $G' - S'$ and $|S'| = s$. We show that $r = s + d + 2$. Otherwise, $r \geq s + d + 3$ and so $r \geq s + d + 4$ by parity. Let $e = c_1 c_2$, where $c_1 \in V(C_1)$ and $c_2 \in V(C_2)$. Clearly, $(G \cup e) - (R \cup M \cup S')$ contains at least $s + d + 2$ odd components, i.e., $G \cup e$ is not an (n, k, d) -graph, a contradiction to the fact that G is a maximal non- (n, k, d) -graph.

We next show that $G' - S'$ has no even components. Otherwise, assume that $G' - S'$ contains an even component D . Let $e = dc_1$, where $d \in D$ and $c_1 \in V(C_1)$, and consider $G \cup e$. Clearly, $(G \cup e) - (R \cup M \cup S')$ contains exactly $s + d + 2$ odd components since the components D and C_1 together with the edge e form an odd component of $G \cup e$. Thus $G \cup e$ is not an (n, k, d) -graph, a contradiction.

Finally we show that $G[R \cup M \cup S']$ is complete. Otherwise, there exist vertices x and y in $R \cup M \cup S'$ such that $e = xy \notin E(G)$. Consider $G \cup e$. Since $(G \cup e) - (R \cup M \cup S')$ contains exactly $s + 2 + d$ odd components, $G \cup e$ is not an (n, k, d) -graph, a contradiction. By a similar argument, it is easy to see that each C_i is complete for $1 \leq i \leq s + d + 2$. Furthermore, each vertex of C_i ($1 \leq i \leq s + d + 2$) is adjacent to every vertex of $G[R \cup M \cup S']$.

Now, for $1 \leq i \leq s + d + 2$, let $|V(C_i)| = 2t_i + 1$ for some non-negative integer t_i . Then $p = |V(G)| = n + 2k + s + \sum_{i=1}^{s+d+2} |V(C_i)| = n + 2k + 2s + d + 2 + 2 \sum_{i=1}^{s+d+2} t_i \geq n + 2k + 2s + d + 2$. Therefore, $0 \leq s \leq \frac{p-n-2k-d}{2} - 1$ and $\sum_{i=1}^{s+d+2} t_i = \frac{p-n-2k-d}{2} - s - 1$ are as required. This completes the proof of the theorem. \square

Acknowledgement

The third author is supported by the Discovery Grant (144073) of the Natural Sciences and Engineering Research Council of Canada.

References

- [1] N. Ananchuen, L. Caccetta, W. Ananchuen, A characterization of maximal non- k -factor-critical graphs, *Discrete Math.* 307 (2007) 108–114.
- [2] C. Berge, Sur le couplage maximum d'un graphe, *C. R. Acad. Sci., Paris* 247 (1958) 258–259.
- [3] J. Bondy, U.S.R. Murty, *Graph Theory with Applications*, The Macmillan Press, London, 1976.
- [4] O. Favaron, On k -factor-critical graphs, *Discuss. Math. Graph Theory* 16 (1996) 41–51.
- [5] Z. Jin, H. Yan, Q. Yu, Generalization of matching extensions in graphs (II), *Discrete Appl. Math.* 155 (2007) 1267–1274.
- [6] G. Liu, Q. Yu, Generalization of matching extensions in graphs, *Discrete Math.* 231 (2001) 311–320.
- [7] G. Liu, Q. Yu, On (n, k) -extendable graphs and induced subgraphs, *Int. Math. Forum* 2 (2007) 1141–1148.
- [8] D. Lou, On matchability of graphs, *Australas. J. Combin.* 21 (2000) 201–210.
- [9] M. Plummer, On n -extendable graphs, *Discrete Math.* 31 (1980) 201–210.
- [10] Q. Yu, Characterizations of various matching extensions in graphs, *Australas. J. Combin.* 7 (1993) 55–64.